

# Prenatal Diagnosis of 30 Fetuses at Risk for Fragile X Syndrome

M. Grasso, L. Perroni, S. Colella, G. Piombo, A. Argusti, M. Lituania, M. Buscaglia, U. Giussani, M.G. Grimoldi, and F. Dagna Bricarelli

*Centro di Genetica Umana, E.O. Ospedali Galliera, Genova (M.G., L.P., S.C., G.P., A.A., F.D.B.); Divisione di Ostetricia e Ginecologia, A.O. Santa Corona, Pietra Ligure (M.L.); Istituto Scienze Biomediche, Ospedale San Paolo, Università degli Studi, Milano (M.B., M.G.G.); SIT, Sezione di Citogenetica, Ospedali Riuniti di Bergamo, Italy (U.G.)*

**The results of 30 prenatal diagnoses for fragile X syndrome are reported. Amniotic fluid cells were examined in 1 case, fetal blood in 4, and chorionic villi samples in the others. Of the 5 fetuses analyzed by cytogenetic methods, 1 had showed 4% of fraXq27.3 expression sites and the pregnancy was terminated. For 1 diagnosis, linkage analysis was used: the female fetus turned out to be normal. In 24 fetuses, the direct analysis of the mutation by StB12.3 probe was performed: 6 female and 3 male fetuses were found to carry a full mutation and 1 female fetus was found to carry a premutation. In 3 cases, the diagnoses were verified on fetal blood samples. Several tissues of 2 aborted male fetuses were analyzed for the fragile X mutation. The results are reported and discussed.** © 1996 Wiley-Liss, Inc.

**KEY WORDS:** prenatal diagnosis, fragile X syndrome

## INTRODUCTION

The availability of methods for the direct detection of the FMR1 mutation has made prenatal diagnosis of fragile X syndrome easy to perform, with the results being more informative than those obtained by cytogenetic testing and linkage analysis.

Direct detection is based on the evaluation of the size and methylation pattern of the FMR1 CGG repeats [Kremer et al., 1991; Oberlé et al., 1991; Verkerk et al., 1991]; early prenatal studies showed no differences in the mutation pattern between amniotic fluid cells

[Dobkin et al., 1991] and leukocytes, and chorionic villi samples (CVS) presented some unique aspects. In CVS, full mutations may have greater size heterogeneity than in leukocytes: normal methylation related to X inactivation is not clearly evident before the 12th week of gestation, and fragile X full mutation may be not hypermethylated [Hirst et al., 1991; Sutherland et al., 1991; Devys et al., 1992; Suthcliffe et al., 1992].

We report our experience with fragile X prenatal diagnosis, which was performed by cytogenetic analysis before 1989 and then by molecular methods.

## MATERIALS AND METHODS

### Patients

From 1982 to June 1995, 30 prenatal diagnoses were performed in 28 women (1 had 2 pregnancies and 1 a twin pregnancy). Twenty-seven of the 28 mothers had 1 or more relatives known to be affected by fragile X syndrome; the remaining mother had a family history positive for undiagnosed mental retardation.

In the 6 mothers whose prenatal diagnoses were performed before the StB12.3 probe was available, it was not possible to determine whether they carried a premutation or a full mutation.

Of the 22 women whose carrier status was ascertained by the direct analysis of the mutation, 11 had a premutation and 11 a full mutation.

### Fetal Samples

For cytogenetic analysis, amniotic cells were used in 1 case and fetal blood in 4. Molecular analysis was carried out on CVS; in 3 fetuses, the results were not straightforward and, for this reason, were confirmed on blood samples. Constitutional karyotype analysis was performed for all cases.

### Cytogenetic Analysis

Constitutional chromosome analysis was performed according to standard techniques; in addition, for the first 5 prenatal diagnoses, the following methods of fragile X induction were used: (1) TC199 + 5% AB serum and (2) TC199 + trimethoprim 15 µg/ml.

Received for publication September 15, 1995; revision received December 1, 1995.

Address reprint requests to Dr. Lucia Perroni, Centro di Genetica Umana, E.O. Ospedali Galliera, Via Volta 10, 16128 Genova, Italy.

TABLE I. Prenatal Diagnoses Performed by Cytogenetic Analysis

Pedigree i.d.	Tissue	Sex of fetus	fraXq27.3
45/85	Amniotic fluid cells	Male	Negative
332/86	Fetal blood	Male	Negative
45/87	Fetal blood	Male	Negative
260/88	Fetal blood	Female	Negative
172/89	Fetal blood	Male	4%

### Molecular Analysis

The CVS DNA was obtained by phenol chlorophorm extraction. Seven micrograms were digested with EcoRI/EagI and with BglII to get sharper smears in cases of faint full mutations. Blotting and hybridization with the probe StB12.3 (kindly provided by Dr. J.L. Mandel) were carried out according to Rousseau et al. [1991]. In the only diagnosis performed by linkage analysis, a number of polymorphic sites flanking the FRAXA locus were investigated.

### RESULTS

All DNA and cytogenetic results are reported in chronological order in Tables I and II.

From 1982 to 1989, fraXq27.3 site induction was used for 5 prenatal diagnoses; 1 male fetus showed 4% of fraXq27.3 expression, and the pregnancy was terminated (Table I).

The 25 prenatal diagnoses performed by molecular methods are listed in Table II.

In case 1, PRS91, haplotype analysis demonstrated that the female fetus had inherited the normal X chromosome from the mother.

Twelve fetuses of 11 premutated women were investigated: 8 were normal (5 female and 3 male), 3 were carriers of a full mutation (1 female and 2 male), and 1 twin female had a premutation.

Among the 12 fetuses of the 11 fully mutated mothers, 6 were normal (5 female and 1 male) and 6 carried a full mutation (5 female and 1 male).

In 3 cases, it was necessary to check the fetal blood results. In case 9, BSL94, we considered it necessary to verify the methylation status of the full mutation on fetal blood because the analysis on CVS of the male fetus showed an unmethylated full mutation ( $\Delta > 800$  bp). DNA analysis, performed at the 23rd week of pregnancy, showed a methylated full mutation.

We also faced problems with the twin pregnancy because, on CVS analysis, 1 of the 2 female fetuses had normal results and 1 was premutated. Because chorionic villi are not fetal tissues, we wanted to rule out the possibility of mosaicism in the premutated fetus. Therefore, we examined the DNA from blood leucocytes of both fetuses, and the diagnoses were confirmed.

All but 2 of the pregnancies where a fetus carried a full mutation were terminated. One fragile X normal pregnancy was terminated because of trisomy 21.

Five pregnancies are still ongoing; follow-up blood samples were obtained on 11 cases predicted to be normal and on 5 predicted to have full mutations. All predictions were confirmed.

TABLE II. Prenatal Diagnoses Performed by Molecular Analysis<sup>a</sup>

Case	Pedigree i.d.	Maternal mutation	CVS (gestational age)	Fetal blood	Sex of fetus	RFLPs	StB12.3
1	PRS (91)	—	14	—	Female	Normal	—
2	1PGG (92)	Full mutation	11	—	Female	N.I. <sup>c</sup>	Normal
3	<b>GRM (92)</b>	<b>Full mutation</b>	<b>11</b>	—	<b>Female</b>	<b>N.I.</b>	<b>Full mutation</b>
4	FNL (92)	Premutation	18	—	Female	—	Normal
5	MNT (93)	Full mutation	10	—	Female	—	Full mutation
6	RND (93)	Premutation	10	—	Female	—	Normal
7	2PGG (94)	Full mutation	11	—	Female	—	Normal
8	FLR (94)	Premutation	11	—	Female	—	Normal
9	<b>BSL (94)</b>	<b>Premutation</b>	<b>18</b>	<b>Yes</b>	<b>Male</b>	—	<b>Full mutation</b>
10	<b>GRP (94)</b>	<b>Full mutation</b>	<b>13</b>	—	<b>Female</b>	—	<b>Full mutation</b>
11	DLC (94)	Full mutation	11	—	Female	—	Normal
12	PNT (94)	Full mutation	11	—	Male	—	Normal
13	<b>BDN (94)</b>	<b>Full mutation</b>	<b>12</b>	—	<b>Male</b>	—	<b>Full mutation</b>
14	PLB (94)	Premutation	12	—	Male	—	Normal
15	LMI (94)	Premutation	14	—	Female	—	Normal
16	<b>PNU (95)</b>	<b>Full mutation</b>	<b>11</b>	—	<b>Female</b>	—	<b>Normal<sup>b</sup></b>
17	BGN (95)	Premutation	12	—	Male	—	Normal
18	RCL (95)	Full mutation	14	—	Female	—	Normal
19	<b>FRT (95)</b>	<b>Premutation</b>	<b>12</b>	—	<b>Male</b>	—	<b>Full mutation</b>
20	BDV (95)	Premutation	11	—	Male	—	Normal
21	<b>RST (95)</b>	<b>Full mutation</b>	<b>12</b>	—	<b>Female</b>	—	<b>Full mutation</b>
22	<b>BVN (95)</b>	<b>Full mutation</b>	<b>12</b>	—	<b>Female</b>	—	<b>Full mutation</b>
23	IPL (95)	Premutation	13	—	Female	—	Full mutation
24	<b>MMN (95)</b>	<b>Premutation</b>	<b>10</b>	<b>Yes</b>	<b>Female</b>	—	<b>Normal</b>
25				Yes	Female		Premutation

<sup>a</sup>Terminated pregnancies are indicated by boldface type.

<sup>b</sup>Pregnancy terminated because of trisomy 21.

<sup>c</sup>NI = not informative.

Of the 2 fully mutated male fetuses, BSL94 and BDN94, it was possible to examine a number of tissues: brain, liver, skin, heart, spleen, kidney, lung, testicles, cartilage, and muscle. The 2 fetuses were of different gestational ages: BSL was 25 weeks and BDN was 15 weeks. BSL's mother was premutated and BDN's was fully mutated. The results are shown in Figs. 1 and 2: in BDN, a mosaic premutation/full mutation was present in all the tissues examined (Fig. 1), and a premutated allele was clearly seen only in the lung of fetus BSL (data not shown). In the cartilage and voluntary muscle of both fetuses, smaller bands of 2.8 kb in BDN and 2.6 kb in BSL (Fig. 2) were also present.

### DISCUSSION

We found that CVS are the tissue of preference for fragile X prenatal diagnosis because of the speed with which the results are obtained. We had no problems due to maternal contamination, and in all but 3 cases, results on CVS were clear enough to not need fetal blood confirmation. BglII digestion provides better resolution because it narrows the faint full mutation smears. Analysis of the tissues of the aborted fully mutated fetuses can provide information about the mechanisms of amplification, especially when the fetuses are at different gestational ages, as with our 2 cases.

Mosaic premutation/full mutation was demonstrated in all examined tissues from a 15-week male fetus whose mother carried a full mutation. One hypothesis to explain this finding is that the maternal gamete carried a premutation and that the amplification was a postzygotic event [Reyniers et al., 1993].




Fig. 2. StB12.3 hybridization pattern of EcoRI/EagI double-digested DNA samples derived from BSL fetal tissues (25 gestational weeks). **Lane 1:** Brain. **Lane 2:** Brain of a normal female fetus. **Lane 3:** Liver. **Lane 4:** Liver of a normal male fetus. **Lane 5:** Voluntary muscle. **Lane 6:** Cartilage. **Lane 7:** Testicle.

The 25-week male fetus whose mother carried a premutation showed a clear pre/full mutation mosaicism only in the lung; presumably, the premutated sequence had already evolved into a full mutation in the other tissues.

It remains to be explained why a smaller allele was clearly present (2.8 kb in 1 fetus and 2.6 in the other) in the cartilage and muscle of both fetuses. The instability of the CGG repeats might be the mechanism that leads to regressions, but whether or not a regression in size in the same tissues of the 2 fetuses is due to chance remains to be clarified.

### ACKNOWLEDGMENTS

We thank Domenico Carratta for technical assistance and Dr. Francesca Faravelli for her helpful collaboration. DNA samples and normal and fragile X fetal tissues are stored at the Galliera Genetic Bank, which is supported by Telethon Italia grant C04.

### REFERENCES

- Devys D, Biancalana V, Rousseau F, Boué J, Mandel JL, Oberlé I (1992): Analysis of full fragile X mutation in fetal tissues and monozygotic twins indicate that abnormal methylation and somatic heterogeneity are established early in development. *Am J Med Genet* 43:208-216.
- Dobkin CS, Ding XH, Jenkins EC, Krawczun MS, Brown WT, Goonewardena P, Willner J, Benson C, Heitz D, Rousseau F (1991): Prenatal diagnosis of fragile X syndrome. *Lancet* 338:957-958.
- Hirst M, Knight S, Devys K, Cross G, Ocraft K, Raeburn S, Heeger S, Eunpu D, Jenkins EC, Lindenbaum R (1991): Prenatal diagnosis of fragile X syndrome. *Lancet* 338:956-957.
- Kremer EJ, Pritchard M, Lynch M, Yu S, Holman K, Baker E, Warren ST, Schlessinger D, Sutherland GR, Richards RI (1991): Mapping




Fig. 1. StB12.3 hybridization pattern of EcoRI/EagI double-digested DNA samples derived from BDN fetal tissues (15 gestational weeks). **Lane 1:** Lung of a normal female fetus. **Lane 2:** Brain. **Lane 3:** Liver. **Lane 4:** Voluntary muscle. **Lane 5:** Cartilage. **Lane 6:** Testicle. **Lane 7:** Heart. **Lane 8:** Lung. **Lane 9:** Spleen. **Lane 10:** Skin.

- of DNA instability at the fragile X to a trinucleotide repeat sequence p(CCG)<sub>n</sub>. *Science* 252:1711–1714.
- Oberlé I, Rousseau F, Heitz D, Krets C, Devys D, Hanauer A, Boué J, Bertheas MF, Mandel JL (1991): Instability of a 550 bp DNA segment and abnormal methylation in fragile X syndrome. *Science* 252:1097–1102.
- Reyniers E, Vits L, De Boulle K, van Roy B, van Belzen D, de Graff E, Verkerk AJ, Jorens H, Darby ZJ, Oostra BA, Willems PJ (1993): The full mutation in the FMR 1 gene of male fragile X patients is absent in their sperm. *Nature Genet* 4:143–146.
- Rousseau F, Heitz D, Biancalana V, Blumenfeld S, Kretz C, Boué J, Tommerup N, van der Hagen C, de Lozier-Blanchet C, Croquette MF, et al. (1991): Direct diagnosis by DNA analysis of the fragile X mental retardation syndrome. *N Engl J Med* 325:1673–1681.
- Suthcliffe JS, Nelson DL, Zhang F, Pieretti M, Caskey CT, Saxe D, Warren ST (1992): DNA methylation represses FMR1 transcription in fragile X syndrome. *Hum Mol Genet* 1:397–400.
- Sutherland GR, Gedeon A, Kornman L, Donnelly A, Byard RW, Mulley JC, Cremer E, Lynch M, Pritchard M, Yu S, Richards RI (1991): Prenatal diagnosis of fragile X syndrome by direct detection of the unstable DNA sequence. *N Engl J Med* 325:1720–1722.
- Verkerk AJMH, Pieretti M, Sutcliffe JS, Fu Y, Kuhl DPA, Pizzuti A, Reiner O, Richards S, Victoria MF, Zhang F, Eussen BE, van Ommen GJB, Blonden LAJ, Riggins GJ, Chastain JL, Kunst CB, Galjaard H, Caskey CT, Nelson DL, Oostra BA, Warren ST (1991): Identification of a gene (FMR-1) containing a CGG repeat coincident with a breakpoint cluster region exhibiting length variation in fragile X syndrome. *Cell* 65:905–914.